

# TECHNICAL DATA SHEET

## TRYPTO-CASEIN SOY (TSA) AGAR

### NUTRIENT MEDIA

#### 1 INTENDED USE

Trypto Casein-Soy Agar is a universal nutrient medium suitable for a wide range of uses. In light of its excellent nutritive value, it can be used for the growth and isolation of both aerobic and anaerobic bacteria and to favor the development of the most fastidious microorganisms. Poured in plates or on strips, it is useful in rapid tests for examining the contamination of surfaces. In addition, under the denomination TSA (Tryptic Soy Agar), this formula corresponds to the reference medium used for the evaluation of productivity and selectivity criteria in the context of the ISO 11133 standard.

The typical composition corresponds to that defined in the US and EU Pharmacopeia.

#### 2 PRINCIPLES

The combination of Tryptone and Papaic digest of soybean meal leads to optimal growth, due to the synergy between the protein supply of casein and the carbohydrate supply of soybeans, favoring the growth of most fastidious or non-fastidious microorganisms.

Sodium chloride maintains osmotic balance.

#### 3 TYPICAL COMPOSITION

The composition can be adjusted in order to obtain optimal performance.

For 1 liter of media :

- Tryptone ..... 15,0 g
- Papaic digest of soybean meal ..... 5,0 g
- Sodium chloride ..... 5,0 g
- Bacteriological agar..... 15,0 g

pH of the ready-to-use media at 25 °C : 7,3 ± 0,2.

#### 4 PREPARATION

##### Preparation from dehydrated media :

- Dissolve 40,0 g of dehydrated media (BK047) in 1 liter of distilled or demineralized water.
- Slowly bring to boiling, stirring with constant agitation until complete dissolution.
- Dispense in tubes or flasks.
- Sterilize in an autoclave at 121°C for 15 minutes.
- Cool and maintain the media in a molten state at 44-47 °C.

✓ **Reconstitution :**  
40,0 g/L

✓ **Sterilization :**  
15 min at 121 °C

##### Use of ready-to-melt media :

- Using the ready-to melt media (BM017, BM049), or if the media was prepared in advance as above, melt the agar for the minimum amount of time necessary to achieve total liquefaction.
- Cool and maintain the media in a molten state at 44-47 °C.

## 5 INSTRUCTIONS FOR USE

### Surface enumeration

- Pour the molten media held at 44-47 °C into sterile Petri plates.
- Let solidify on a cold, flat surface.
- Dry the plates in an incubator with the covers partially removed.
- To the surface of plates prepared as above or using pre-poured plates (BM050), inoculate 0.1 mL of the test sample and its serial with a sterile triangle or "hockey stock".
- Incubate according to the reference applied :
  - at 30-35 °C for 72 ± 6 hours (NF EN ISO 21149, NF EN ISO 18415)
  - at 30-35 °C for 3 to 5 days for Total viable count (Pharmacopeia)

✓ **Inoculation :**  
0,1 mL on surface

✓ **Incubation :**  
30-35 °C

### NOTE :

The media can also be used after having filtered through a nitrocellulose membrane (Pharmacopeia, NF EN ISO 21149)

### Enumeration in pour plates

- Transfer 1 mL of the initial suspension and its serial dilutions to successive sterile Petri plates. .
- Pour roughly 15 mL of molten media per plate.
- Homogenize well and let solidify on a cold, flat surface.
- Incubate according to the reference applied :
  - at 30-35 °C for 72 ± 6 hours (NF EN ISO 21149, NF EN ISO 18415)
  - at 30-35 °C for 3 to 5 days for Total Aerobic Viable Count (Pharmacopeia)

✓ **Inoculation :**  
1 mL in pour plates

✓ **Incubation :**  
30-35 °C

For other uses, refer to the references in vigor.

## 6 RESULTS

Following incubation, observe bacterial growth.

When performing enumeration, only retain the plates containing between 30 and 300 colonies.

## 7 QUALITY CONTROL

**Dehydrated media :** cream-white powder, free-flowing and homogeneous.

**Prepared media :** amber agar.

Typical culture response after incubation 48 h at 30-35 °C (NF EN ISO 11133) :

Microorganisms		Growth (Productivity Ratio : $P_R$ )
<i>Staphylococcus aureus</i>	WDCM 00032	$P_R \geq 50 \%$
<i>Pseudomonas aeruginosa</i>	WDCM 00026	$P_R \geq 50 \%$
<i>Bacillus cereus</i>	WDCM 00001	$P_R \geq 70 \%$
<i>Bacillus subtilis</i> ssp. <i>spizizenii</i>	WDCM 00003	$P_R \geq 70 \%$
<i>Escherichia coli</i>	WDCM 00012	$P_R \geq 70 \%$
<i>Listeria monocytogenes</i> 4 b	WDCM 00021	$P_R \geq 70 \%$
<i>Staphylococcus aureus</i>	WDCM 00034	$P_R \geq 70 \%$

## 8 STORAGE / SHELF LIFE

---

**Dehydrated media** : 2-30 °C.

**Ready-to-melt media in vials** : 2-25 °C

**Pre-poured media in Petri plates** : 2-8 °C

The expiration dates are indicated on the labels.

**Prepared media in tubes or vials (\*)** : 180 days at 2-25 °C.

**Prepared media in plates (\*)** : 30 days at 2-8 °C.

(\*) Benchmark value determined under standard preparation conditions, following manufacturer's instructions.

## 9 PACKAGING

---

**Dehydrated media :**

500 g bottle ..... BK047HA

**Pre-poured media in Petri plates (Ø 90 mm) :**

20 plates ..... BM05008

**Ready-to-melt media in vials :**

10 x 100 mL vials ..... BM01708

10 x 200 mL vials ..... BM04908

## 10 BIBLIOGRAPHY

---

NF EN ISO 9308-1. Septembre 2000. Qualité de l'eau. Recherche et dénombrement des *Escherichia coli* et des bactéries coliformes. Partie 1 : Méthode par filtration sur membrane.

NF EN 15784. Décembre 2009. Aliments des animaux. Isolement et dénombrement des souches de *Bacillus* spp. présumées.

NF EN ISO 11930. Juin 2012. Cosmétiques - Microbiologie - Évaluation de la protection antimicrobienne d'un produit cosmétique.

NF EN ISO 11133. Juillet 2014. Microbiologie des aliments, des aliments pour animaux et de l'eau - Préparation, production, stockage et essais de performance des milieux de culture (Tirage 2 (2016-01-01)).

NF EN ISO 21150. Février 2016. Cosmétiques. Microbiologie. Détection d'*Escherichia coli*.

NF EN ISO 22717. Février 2016. Cosmétiques. Microbiologie. Recherche de *Pseudomonas aeruginosa*.

NF EN ISO 22718. Février 2016. Cosmétiques. Microbiologie. Détection de *Staphylococcus aureus*.

ISO 9308-1. Septembre 2014. Qualité de l'eau - Dénombrement des *Escherichia coli* et des bactéries coliformes - Partie 1 : méthode par filtration sur membrane pour les eaux à faible teneur en bactéries.

NF EN ISO 22964. Juin 2017. Microbiologie de la chaîne alimentaire - Méthode horizontale pour la détection de *Cronobacter* spp..

NF EN ISO 11731. Juillet 2017. Qualité de l'eau. Dénombrement des *Legionella*.

NF EN ISO 18415. Août 2017. Cosmétiques. Microbiologie. Détection des micro-organismes spécifiés et non spécifiés.

NF EN ISO 21149. Août 2017. Cosmétiques. Microbiologie. Dénombrement et détection des bactéries aérobies mésophiles.

Pharmacopée Européenne. Chapitre 2.6.13. Contrôle microbiologique des produits non stériles : Recherche de microorganismes spécifiés.

## 11 ADDITIONAL INFORMATION

---

The information provided on the labels take precedence over the formulations or instructions described in this document and are susceptible to modification at any time, without warning.

Document code : TRYPTO CASEINE SOJA AGAR\_EN v11

Creation date : 06-2003

Updated : 01-2018  
Origin of revision : Bibliography.